

Explanatory Notes

All cases (locked and unlocked) admitted to hospital between 01 November 2025 and 31 January 2026 have been included. Only cases where the necessary data are available have been included in the denominator for each individual analysis.

At hospital level, runcharts are compared to hospitals within the same ICB.

The results for process measures for which fewer than 10 cases have available data will not be reported. Instead the value will be marked as 'Insufficient data'.

Mortality

This section defines three key mortality measures for the monthly report. In all cases we include only patients whose surgery-to-discharge interval (Q4.1 - Q7.8) is ≤ 30 days, and we exclude any with missing discharge status (Q7.7) or missing dates (Q4.1/Q7.8).

1. 30-Day Observed (Crude) Mortality Rate

Let

- d = number of patients who **died** within 30 days of surgery,
- N = total number of patients with known discharge status (alive, died, or still in hospital at 60 days).

Then the crude 30-day mortality rate (as a percentage) is

$$\text{Crude 30-day Mortality Rate} = \frac{d}{N} \times 100.$$

2. Standardised Mortality Ratio (SMR)

Let

- $O = d$ = observed deaths within 30 days,
- $E = \sum_i \text{RiskScore}_i$ = sum of individual parsimonious NELA mortality risk scores for all N patients.

The SMR is

$$\text{SMR} = \frac{O}{E}.$$

3. Risk-Adjusted Mortality

Combines the SMR with the **National** 30-day mortality rate for the examined three month period:

$$\text{Risk-Adjusted Mortality} = \text{SMR} \times (\text{National 30-day mortality}) \times 100.$$

For better insight to how these standards have been structured, please refer to the **NELA standards document**.

Quarterly mean performance



Overall performance



Risk-adjusted mortality

Rating boundaries are lower and upper 99.8% and 95% confidence limits



Non-risk-adjusted measures

Rating boundaries are lower and upper national quartiles




Leicester Royal Infirmary

2025-26 Reporting Period 10: 01 November 2025 - 31 January 2026

These plots represent patients having an emergency laparotomy during Year 2025-26 Reporting Period 10 of NELA data collection. This version will be made publicly available via the NELA website. Feedback from participating hospitals is welcome.



NELA process and outcome measures



Estimated case ascertainment
01 November 2025 - 31 January 2026

**Estimated case ascertainment
(Based on HES/PEDW Data)**



Expected number of cases 105
Total cases entered 14
Cases locked 6
Cases unlocked 8



Proportion of patients who had a CT scan that was reported by a senior radiologist (ST3+) and communicated with the team in the correct time scale before surgery
01 November 2025 - 31 January 2026

CT reported by a senior radiologist (ST3+) and communicated with the team in the correct time scale before surgery.



National mean 11%
ICB mean 0%
Number of patients included 8
Data completeness 100%



Proportion of patients with suspected sepsis or infection that have antibiotic administration within the correct clinical timeframe
01 November 2025 - 31 January 2026



Combined Infection management standard - antibiotic administration within the correct clinical timeframe

National mean 20%
ICB mean 33%
Number of patients included 6
Data completeness 55%





Sepsis/septic shock - antibiotic administration within the correct clinical timeframe

National mean 12%
ICB mean 25%
Number of patients included 4
Data completeness 44%





Infection - antibiotic administration within the correct clinical timeframe

National mean 28%
ICB mean 50%
Number of patients included 2
Data completeness 18%





Non-elective, immediate surgery cases, arriving in theatre within 6 hours.

National mean 12%
ICB mean 0%
Number of patients included 7
Data completeness 100%





Risk documented before surgery

National mean 75%
ICB mean 29%
Number of patients included 14
Data completeness 100%





Risk documented after surgery

National mean 55%
ICB mean 0%
Number of patients included 14
Data completeness 100%





Admitted to Critical Care (risk of death \geq 5%)

National mean 63%
ICB mean 33%
Number of patients included 12
Data completeness 100%





Consultant Anaesthetist & Consultant Surgeon in theatre (risk of death \geq 5%)

National mean 80%
ICB mean 50%
Number of patients included 12
Data completeness 100%





Perioperative Assessment by a member of the geriatrician-led multidisciplinary team for patient aged 65 or over and frail (CFS \geq 5) or 80+

National mean 31%
ICB mean 0%
Number of patients included 4
Data completeness 44%



Median postoperative length of stay


National median 10 days
ICB median 8 days
Number of patients included 6
Data completeness 86%



Mean postoperative length of stay

National mean 14 days
ICB mean 10 days
Number of patients included 6
Data completeness 86%

Risk-Adjusted Mortality



Number of patients included 6 | 30-day risk-adjusted mortality rate NaN% | National 30-day mortality rate 7.6%

Integrated Care Board

Leicester Royal Infirmary is part of the NHS Leicester, Leicestershire And Rutland Integrated Care Board ICB. This comprises Leicester Royal Infirmary, Leicester General Hospital, Glenfield Hospital.