

Explanatory Notes

All cases admitted to hospital between 01 February 2025 and 30 April 2025 have been included. Only cases where the necessary data are available have been included in the denominator for each individual analysis.

The results for process measures for which fewer than 10 cases have available data will not be reported. Instead value will be marked as 'Insufficient data'.

For this report runcharts are compared to national quartiles.

For better insight to how these standards have been structured, please refer to the **NELA standards document**.




NHS Lancashire And South Cumbria Integrated Care Board

2025-26 Reporting Period 1: 01 February 2025 - 30 April 2025

These plots represent patients having an emergency laparotomy during Year 2025-26 Reporting Period 1 of NELA data collection.



NELA process and outcome measures



Estimated case ascertainment
01 February 2025 - 30 April 2025

**Estimated Case Ascertainment
(Based on HES Data)**



Expected number of cases 192
Total cases entered 154
Cases locked 131
Cases unlocked 23



Proportion of patients who had a CT scan that was reported by senior radiologist and communicated with the team in the correct time scale before surgery
01 February 2025 - 30 April 2025

CT reported by a senior radiologist (ST3+) and communicated with the team in the correct time scale before surgery.



National mean 13%
Number of patients included 116
Data completeness 100%



Proportion of patients with suspected sepsis/septic shock or infection that have antibiotic administration within the correct clinical timeframe
01 February 2025 - 30 April 2025



Combined Infection management standard - antibiotic administration within the correct clinical timeframe

National mean 21%
Number of patients included 62
Data completeness 97%





Sepsis/septic shock - antibiotic administration within the correct clinical timeframe

National mean 13%
Number of patients included 31
Data completeness 93%





Infection - antibiotic administration within the correct clinical timeframe

National mean 31%
Number of patients included 52
Data completeness 85%





Timeliness - proportion of patients arriving in theatre according to correct clinical timeframe.

National mean 11%
Number of patients included 90
Data completeness 100%



Risk documented before surgery



National mean 76%
Number of patients included 154
Data completeness 100%



Risk of death documented after surgery
01 February 2025 - 30 April 2025

Risk documented after surgery



National mean 59%
Number of patients included 154
Data completeness 100%



Admitted to critical care following surgery when the risk of death \geq 5% (Excludes patients who died in theatre or with a decision to palliate)
01 February 2025 - 30 April 2025

Admitted to Critical Care (risk of death \geq 5%)



National mean 65%
Number of patients included 83
Data completeness 100%



Consultant surgeon and anaesthetist present in theatre when risk of death \geq 5%
01 February 2025 - 30 April 2025

Consultant Anaesthetist & Consultant Surgeon in theatre (risk of death \geq 5%)



National mean 80%
Number of patients included 84
Data completeness 82%



Perioperative assessment by a care of the older person specialist
01 February 2025 - 30 April 2025



Perioperative Assessment by a member of the geriatrician-led multidisciplinary team for patient aged 65 or over and frail (CFS \geq 5) or 80+

National mean 34%
Number of patients included 44
Data completeness 98%



Median postoperative Length of Stay


National median 10 days
Number of patients included 119
Data completeness 100%



Mean postoperative Length of Stay

National mean 14 days
Number of patients included 119
Data completeness 100%

Risk-Adjusted Mortality



Number of patients included 134 | 30-day risk-adjusted mortality rate 6.8% | National 30-day mortality rate 7.8%